

TABLE 2. *Methods used in retrospective and cross sectional studies of peptic ulcer and smoking (cont.)*

Author, year, country, reference	Sex	Number	Method of selection	Controls		Comments
				Number	Method of selection	
Kasanen and Forsström, 1966, Finland (13).	M	43 Gastric. 57 Duodenal.	Successive male admissions with peptic ulcer treated at medical clinic or outpatient department of University Hospital. Only patients under 65 years of age or those who had been working were included.	100	Successive men treated at medical clinic who had no gastrointestinal symptoms or signs of CHD.	A special questionnaire was used for the interview.
Gillies and Skyring, 1968, 1968, Australia (9).	M and F	100 Gastric. 50 Duodenal.	Patients with peptic ulcer were selected from hospital admissions in 1967.	150	Matched by age and sex from the same ward at the same time and with absence of signs or symptoms or past history of upper gastrointestinal disease.	Diagnosis well established with X-ray, gastroscopy, or surgery.
Gillies and Skyring, 1969, Australia (10).	M and F	10 Gastric. 48 Duodenal. 18 Uncertain location.	1,405 workers from a broadcasting company, a manufacturing company, and a bus company were interviewed for a history of peptic ulcer.	100 1,329	Two control groups: 1. 100 peptic ulcer patients previously reported by authors. 2. 1,329 workers without ulcer.	All information obtained by question card. All ulcers were proved by X-ray or surgery.
Monson, 1970, U.S.A. (15).	M and F	52 Gastric. 452 Duodenal. 139 Not specified.	643 physicians from Massachusetts who responded affirmatively to a questionnaire sent to them in 1967 asking how many had had a peptic ulcer.	625	Controls were physicians without ulcer disease who were matched to ulcer patients by year of birth.	Diagnosis established by X-ray or surgery except for 46 "clinical" cases.

TABLE 3.—Summary of results of retrospective and cross sectional studies of peptic ulcer and smoking

Author, year, country, reference	Percent nonsmoker		Amount of tobacco used	
	Cases	Controls	Cases	Controls
Barnett, 1927, U.S.A. (2).	Total 18.0 Gastric 15.0 Duodenal 20.0	25.0		
Trowell, 1934, England (21).	Duodenal 8.0	17.0	Average number: Cigarettes .. 12.0 per day.....11.1 per day Pipe 1.6 ounces per week.. 2.15 ounces per week	
Allibone and Flint, 1958, England (1).	38.0	54.0		
Doll et al., 1958, England (7).	Gastric: Males 1.3 Females ... 51.1 Duodenal: Males 2.1 Females ... 53.7	4.7 66.8 5.8 62.0	Gastric: <i>Percent smoking >25 cigarettes per day</i> Males 10.6 Females 1.1 Duodenal: Males 10.2 Females 1.9	11.3 1.1 12.7 1.9
Edwards et al., 1959, England (8).	Percent of peptic ulcer by smoking category			
	Never smoked 6.0			
	Formerly smoked 6.7			
	Cigarettes:			
	1-9 per day 9.4			
	10-19 per day 9.8			
	>20 per day 12.0			
	Pipe 6.5			
	Pipe and cigarettes 8.5			

TABLE 3.—Summary of results of retrospective and cross sectional studies of peptic ulcer and smoking (cont.)

Author, year, country, reference	Percent nonsmoker		Amount of tobacco used	
	Cases	Controls	Cases	Controls
Kasanen and Forsström, 1966, Finland (13).	"Peptic"	10.0	40.0	
			Cigarettes per day:	
			<10	10.0
			10-20	19.0
			20	42.0
			>20	19.0
Gillies and Skyring, 1968, Australia (9).	Gastric	18.0	44.0	
	Duodenal	62.0	71.0	
			Mean number cigarettes per day:	
			Gastric	23.3
			Duodenal ..	23.2
Gillies and Skyring, 1969, Australia (10).			Duration of smoking (years):	
			Gastric	30.2
			Duodenal ..	24.2
Monson, 1970, U.S.A. (15).	Gastric	17.9	55.6	
	Duodenal	36.6		
	Duodenal	32.1	46.7	
	Gastric	19.2		
	Not Specified ..	43.2		
			Percent smoking >20 cigarettes per day	
			Age:	Gastric Duodenal
			20	38.8 27.3 30.1
			30	45.7 43.0 47.1
			45	60.2 49.5 46.9
			60	54.1 40.4 44.0

REFERENCES

- (1) ALLIBONE, A., FLINT, F. J. Bronchitis, aspirin, smoking, and other factors in the aetiology of peptic ulcer. *Lancet* 2: 179-182, July 26, 1958.
- (2) BARNETT, C. W. Tobacco smoking as a factor in the production of peptic ulcer and gastric neurosis. *Boston Medical and Surgical Journal* 197(12): 457-459, September 22, 1927.
- (3) BATTERMAN, R. C., EHRENFELD, I. The influence of smoking upon the management of the peptic ulcer patient. *Gastroenterology* 12(4): 575-585, April 1949.
- (4) COOPER, P., KNIGHT, J. B., JR. Effect of cigarette smoking on gastric secretions of patients with duodenal ulcer. *New England Journal of Medicine* 255(1): 17-21, July 5, 1956.
- (5) DOLL, R., HILL, A. B. Mortality in relation to smoking: 10 years' observations of British doctors. Part I. *British Medical Journal* 1(5395): 1399-1410, May 30, 1964.
- (6) DOLL, R., HILL, A. B. Mortality in relation to smoking: 10 years' observations of British doctors (concluded). *British Medical Journal* 1(5396): 1460-1467, June 6, 1964.
- (7) DOLL, R., JONES, F. A., PYGOTT, F. Effect of smoking on the production and maintenance of gastric and duodenal ulcers. *Lancet* 1: 657-662, March 29, 1958.
- (8) EDWARDS, F., MCKEOWN, T., WHITFIELD, A. G. W. Association between smoking and disease in men over sixty. *Lancet* 1: 196-201, January 24, 1959.
- (9) GILLIES, M., SKYRING, A. Gastric ulcer, duodenal ulcer and gastric carcinoma: A case-control study of certain social and environmental factors. *Medical Journal of Australia* 2(25): 1132-1136, December 21, 1968.
- (10) GILLIES, M. A., SKYRING, A. Gastric and duodenal ulcer. The association between aspirin ingestion, smoking and family history of ulcer. *Medical Journal of Australia* 2(6): 280-285, August 9, 1969.
- (11) HAMMOND, E. C. Smoking in relation to the death rates of 1 million men and women. IN: Haenszel, W. (Editor). *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966. pp. 127-204.
- (12) KAHN, H. A. The Dorn study of smoking and mortality among U.S. veterans: Report on 8½ years of observation. IN: Haenszel, W. (Editor). *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966. pp. 1-125.
- (13) KASANEN, A., FORSSTROM, J. Social stress and living habits in the etiology of peptic ulcer. *Annales Medicinae Internal Fenniae* 55(1): 13-22, 1966.
- (14) MITTY, W. F., JR., ROUSSELOT, L. M., DELANY, G. J. Smoking and its relation to nutritional status of patients following gastrectomy; a five-year follow-up survey of 171 patients. *Annals of Surgery* 150(1): 76-84, July 1959.
- (15) MONSON, R. R. Cigarette smoking and body form in peptic ulcer. *Gastroenterology* 58(3): 337-344, March 1970.

- (16) PACKARD, R. S. Smoking and the alimentary tract: A review. *Gut* 1: 171-174, 1960.
- (17) PALMER, W. L. Peptic ulcer. Chapter 32. IN: Paulson, M. (Editor). *Gastroenterologic Medicine*. Philadelphia, Lea & Febiger, 1969, pp. 710-757.
- (18) SCHNEDORF, J. G., IVY, A. C. The effect of tobacco smoking on the alimentary tract. An experimental study of man and animals. *Journal of the American Medical Association* 112(10): 898-904, March 11, 1939.
- (19) STEIGMANN, F., DOLEHIDE, R. H., KAMINSKI, L. Effects of tobacco smoking on gastric acidity and motility of hospital controls and patients with peptic ulcer. *American Journal of Gastroenterology* 22: 399-409, 1954.
- (20) THOMPSON, J. H. Effects of nicotine and tobacco smoke on gastric secretion in rats with gastric fistulas. *American Journal of Digestive Diseases* 15(3): 209-217, March 1970.
- (21) TROWELL, O. A. The relation of tobacco smoking to the incidence of chronic duodenal ulcer. *Lancet* 1: 808-809, April 14, 1934.
- (22) U.S. PUBLIC HEALTH SERVICE. NATIONAL CENTER FOR HEALTH STATISTICS. *Vital Statistics of the United States—1967. Vol. II—Mortality, Part A*. Washington, U.S. Department of Health, Education and Welfare, Public Health Service Publication, 1969.
- (23) WEIR, J. M., DUNN, J. E., JR. Smoking and mortality: A prospective study. *Cancer* 25(1): 105-112, January 1970.

CHAPTER 7

Tobacco Amblyopia

Contents

	<i>Page</i>
Summary and conclusions	436
References	436

TOBACCO AMBLYOPIA

Tobacco amblyopia (tobacco-alcohol amblyopia) is that syndrome of visual failure occurring in association with the use of tobacco, with or without the concurrent use of alcohol, and with or without concurrent nutritional deficits. The disease has a subacute onset, leading to a loss of visual acuity and color perception (12). It is characterized by centrocecal scotomas which are bilateral but not necessarily symmetrical and which have sloping diffuse edges and by the presence of nuclei of denser visual loss within the large scotomas (22, 23). Such visual impairment is not unique to tobacco amblyopia, as it is also seen in neurodegenerative disorders, such as Leber's hereditary optic atrophy (7, 25).

Clinical information on tobacco amblyopia has appeared in numerous articles throughout the past century. This information has been reviewed by Silvette, et al. (17) and, more recently, by Dunphy (5). Pure tobacco amblyopia (TA), that is amblyopia unassociated with excessive alcohol intake or the exposure to other toxins, is rarely seen in the United States today (12). Walsh, et al. (23) have observed that when TA is found it is usually present in association with nutritional or idiopathic vitamin deficiencies. Victor (22) recently observed that the type of visual defect seen in tobacco amblyopia may be found in clinical circumstances in which tobacco is clearly not a causative factor. He questions whether TA is distinguishable from other forms of amblyopia.

The prevalence of this disorder has been variously estimated in the past at from 0.5 to 1.5 percent of all eye clinic patients (20, 23). However, currently in the United States, it appears to be a rare condition. Silvette, et al. (17) have observed that the incidence of tobacco amblyopia appears to have decreased substantially during the past decades. Other authors (3, 15) have also commented on this trend. Although reference has been made to the increased frequency of certain types of tobacco usage in patients with this disorder, adequate population studies with proper controls have yet to be performed. The association of this disorder with the use of tobacco is strengthened by the frequent clinical observations of improvement following the cessation of smoking although improvement has been noted by some to occur without cessation.

Research into the pathogenesis of tobacco amblyopia has cen-

tered upon the interrelationships of cyanide metabolism, vitamin B₁₂, and other vitamin deficiencies. Three reviews of this material have recently appeared (1, 12, 22). Numerous studies reviewed in these articles suggest that tobacco amblyopia may result from the incomplete detoxification of the cyanide present in tobacco smoke. This failure of detoxification may stem from or be intensified by inadequate dietary intake of necessary nutritional factors. This may be the reason for the association of this disorder with excessive alcohol intake and with its related nutritional deficits (2, 4, 6, 8, 9, 10, 11, 13, 14, 16, 18, 19, 21, 24, 26, 27, 28).

SUMMARY AND CONCLUSIONS

Tobacco amblyopia is presently a rare disorder in the United States. The evidence suggests that this disorder is related to nutritional or idiopathic deficiencies in certain detoxification mechanisms, particularly in handling the cyanide component of tobacco smoke.

REFERENCES

- (1) CANADIAN MEDICAL ASSOCIATION JOURNAL. Tobacco amblyopia. (Editorial) Canadian Medical Association Journal 102(4): 420, February 28, 1970.
- (2) CHISHOLM, I. A., BRONTE-STEWART, J., FOULDS, W. S. Hydroxocobalamin versus cyanocobalamin in the treatment of tobacco amblyopia. Lancet 2(7513): 450-451, August 26, 1967.
- (3) DARBY, P. W., WILSON, J. Cyanide, smoking, and tobacco amblyopia. Observations on the cyanide content of tobacco smoke. British Journal of Ophthalmology 51(5): 336-338, May 1967.
- (4) DREYFUS, P. M. Blood transketolase levels in tobacco-alcohol amblyopia. Archives of Ophthalmology 74(5): 617-620, November 1965.
- (5) DUNPHY, E. B. Alcohol and tobacco amblyopia: A historical survey. American Journal of Ophthalmology 68(4): 569-578, October 1969.
- (6) FOULDS, W. S., BRONTE-STEWART, J. M., CHISHOLM, I. A. Serum thiocyanate concentrations in tobacco amblyopia. Nature 218(5141): 586, May 11, 1968.
- (7) FOULDS, W. S., CANT, J. S., CHISHOLM, I. A., BRONTE-STEWART, J., WILSON, J. Hydroxocobalamin in the treatment of Leber's hereditary optic atrophy. Lancet 1(7548): 896-897, April 27, 1968.
- (8) FOULDS, W. S., CHISHOLM, I. A., BRONTE-STEWART, J., WILSON, T. M. Vitamin B₁₂ absorption in tobacco amblyopia. British Journal of Ophthalmology 53(6): 393-397, June 1969.
- (9) FOULDS, W. S., CHISHOLM, I. A., BRONTE-STEWART, J., WILSON, T. M. The optic neuropathy of pernicious anemia. Archives of Ophthalmology 82(4): 427-432, October 1969.
- (10) FREEMAN, A. G., HEATON, J. M. The aetiology of retrobulbar neuritis in Addisonian pernicious anaemia. Lancet 1(7183): 908-911, April 29, 1961.

- (11) HEATON, J. M., MCCORMICK, A. J. A., FREEMAN, A. G. Tobacco amblyopia: A clinical manifestation of vitamin-B₁₂ deficiency. *Lancet* 2(7041): 286-290, August 9, 1958.
- (12) KNOX, D. L. Neuro-ophthalmology. *Archives of Ophthalmology* 83(1): 103-127, January 1970.
- (13) LINDSTRAND, K., WILSON, J., MATTHEWS, D. M. Chromatography and microbiological assay of vitamin B₁₂ in smokers. *British Medical Journal* 2 (5520): 988-990, October 22, 1966.
- (14) LINNELL, J. C., SMITH, A. D. M., SMITH, C. L., WILSON, J., MATTHEWS, D. M. Effects of smoking on metabolism and excretion of vitamin B₁₂. *British Medical Journal* 2(5599): 215-216, April 27, 1968.
- (15) SCHEPENS, C. L. Is tobacco amblyopia a deficiency disease- Transactions of the Ophthalmological Society of the United Kingdom 66: 309-331, 1946.
- (16) SCHIEVELBEIN, H., WERLE, E., SCHULZ, E. K., BAUMEISTER, R. The influence of tobacco smoke and nicotine on thiocyanate metabolism. *Naunyn-Schmiedeberg's Archiv für Pharmakologie und Experimentelle Pathologie* 262(3): 358-365, February 5, 1969.
- (17) SILVETTE, H., HAAG, H. B., LARSON, P. S. Tobacco amblyopia. The evolution and natural history of a "tobaccogenic" disease. *American Journal of Ophthalmology* 50(1): 71-100, January 1960.
- (18) SMITH, A. D. M. Retrobulbar neuritis in Addisonian pernicious anaemia. (Letter) *Lancet* 1(7184): 1001-1002, May 6, 1961.
- (19) SMITH, A. D. M., DUCKETT, S. Cyanide, vitamin B₁₂, experimental demyelination and tobacco amblyopia. *British Journal of Experimental Pathology* 46(6): 615-622, December 1965.
- (20) TRAQUAIR, H. M. Toxic amblyopia, including retrobulbar neuritis. Transactions of the Ophthalmological Society of the United Kingdom 50: 351-385, 1930.
- (21) VICTOR, M. Tobacco-alcohol amblyopia. A critique of current concepts of this disorder, with special reference to the role of nutritional deficiency in its causation. *Archives of Ophthalmology* 70(3): 313-318, September 1963.
- (22) VICTOR, M. Tobacco amblyopia, cyanide poisoning and vitamin B₁₂ deficiency. A critique of current concepts. Chapter 3. IN: Smith, J. L. (Editor) *Neuro-Ophthalmology. Symposium of the University of Miami and the Bascom Palmer Eye Institute. Hallandale, Florida, Huffman Publishing Co., 1970. pp. 33-48.*
- (23) WALSH, F. B., HOYT, W. F. (Editors) Neurotoxic substances affecting the visual and ocular motor systems. Chapter 15: IN: *Clinical Neuro-Ophthalmology, Volume 3, 3rd Edition. Baltimore, The Williams & Wilkins Company, 1969. pp. 2613-2616.*
- (24) WATSON-WILLIAMS, E. J., BOTTOMLEY, A. C., AINLEY, R. G., PHILLIPS, C. I. Absorption of vitamin B₁₂ in tobacco amblyopia. *British Journal of Ophthalmology* 53(8): 549-552, August 1969.
- (25) WILSON, J. Leber's hereditary optic atrophy: A possible defect of cyanide metabolism. *Clinical Science* 29(3): 505-515, December 1965.
- (26) WILSON, J., MATTHEWS, D. M. Metabolic inter-relationships between cyanide, thiocyanate and vitamin B₁₂ in smokers and nonsmokers. *Clinical Science* 31(1): 1-7, January 1966.

- (27) WOKES, F., PICARD, C. W. The role of vitamin B₁₂ in human nutrition. Clinical Nutrition 3(5) : 383-390, September-October 1955.
- (28) WYNDER, E. L., HOFFMANN, D. Certain constituents of tobacco products. Chapter 8. K. Vapor phase of tobacco smoke. IN: Wynder, E. L., Hoffmann, D. (Editors). Tobacco and Tobacco Smoke. Studies in Experimental Carcinogenesis. New York, Academic Press, 1967. pp. 451-453.